arXiv:1612.05940v1 [math-ph] 18 Dec 2016

PERIODIC AND WEAKLY PERIODIC GROUND STATES FOR THE λ-MODEL ON CAYLEY TREE

FARRUKH MUKHAMEDOV, CHIN HEE PAH, MUZAFFAR RAHMATULLAEV, AND HAKIM JAMIL

Abstract. In this paper we consider the λ-model on the Cayley tree of order two. We describe periodic and weakly periodic ground states for the considered model.

1. Introduction

Each Gibbs measure is associated with a single phase of a physical system [18]. Existence of two or more Gibbs measures means that phase transitions exist. One of fundamental problems is to describe the extreme Gibbs measures corresponding to a given Hamiltonian.

As is known, the phase diagram of Gibbs measures for a Hamiltonian is close to the phase diagram of isolated (stable) ground states of this Hamiltonian. At low temperatures, a periodic ground state corresponds to a periodic Gibbs measure, see [6], [18]. The problem naturally arises on description of periodic and weakly periodic ground states. For the Potts model with competing interactions on the Cayley tree of order two, periodic ground states were studied in [3, 1, 9] (see also [15]). The notion of a weakly periodic ground state was introduced in [17]. For the Ising model with competing interactions, weakly periodic ground states were described in [11], [17]. For the Potts model, such states for normal subgroups of index 2 were studied in [12]. For the Potts model, such states for normal subgroups of index 4 were studied in [13].

On the other hand, it is natural to consider a model which is more complicated than Potts one, therefore, in [7] it was proposed to study so-called λ-model on the Cayley tree (see also [14, 15]). In that model, many possible interactions (nearest-neighbor) are taken into account. In the mentioned paper, for a special kind of λ-model, its disordered phase has been studied (see [8]) and some its algebraic properties were investigated. Furthermore, in [10] we have described ground states of the λ-model on the Cayley tree of order two. This model is much more general than Potts model, and exhibits interesting structure of ground states.

In the present paper, we continue an investigation of the λ-model on the Cayley tree of order two and study periodic and weakly periodic ground states for normal subgroups of index 2. The paper is organized as follows. In Section 2, we recall the main definitions and known facts. In Section 3, we study periodic and weakly periodic ground states. Note that our results extend known ones [16, 17]. Hence, the proposed model opens new perspectives in the theory of statistical mechanics over trees.

2. Preliminaries

Let $\tau^k = (V, L)$ be a Cayley tree of order k, i.e, an infinite tree such that exactly $k+1$ edges are incident to each vertex. Here V is the set of vertices and L is the set of edges of τ^k.

Let G_k denote the free product of $k+1$ cyclic groups $\{e, a_i\}$ of order 2 with generators $a_1, a_2, \ldots, a_{k+1}$, i.e., let $a_i^2 = e$ (see[4]).

There exists a one-to-one correspondence between the set V of vertices of the Cayley tree of order k and the group G_k, see[2],[15].

We show how to construct this correspondence. We choose an arbitrary vertex $x_0 \in V$and associate it with the identity element e of the group G_k. Since we may assume that the graph under consideration is planar, we associate each neighbor of x_0 (i.e., e) with a single generator $a_i, i = 1, 2, \ldots, k+1$, where the order corresponds to the positive direction, see Figure 1.



FIGURE 1. The Cayley tree τ^2 and elements of the group representation of vertices

For every neighbor of a_i, we introduce words of the form a_ia_j . Since one of the neighbors of a_i is e, we put $a_ia_i = e$. The remaining neighbors of a_i are labeled according to the above order. For every neighbor of a_ia_j , we introduce words of length 3 in a similar way. Since one of the neighbors of a_ia_j is a_i, we put $a_ia_ja_j = a_i$. The remaining neighbors of a_ia_j are labeled by words of the form $a_ia_ja_l$, where $i, j, l = 1, 2, ..., k+1$, according to the above procedure. This agrees with the previous stage because $a_ia_ja_j = a_ia_j^2 = a_i$. Continuing this process, we obtain a one-to-one correspondence between the vertex set of the Cayley tree τ^k and the group G_k.

The representation constructed above is said to be *right* because, for all adjacent vertices x and y and the corresponding elements $g, h \in G_k$, we have either $g = ha_i$ or $h = ga_j$ for suitable i and j. The definition of the $left$ representation is similar.

For the group G_k (or the corresponding Cayley tree), we consider the left (right) shifts. For $g \in G_k$, we put

$$T_g(h) = gh \ (T_g(h) = hg) \ for \ all \ h \in G^k$$

The group of all left (right) shifts on G_k is isomorphic to the group G_k.

Each transformation S on the group G_k induces a transformation S on the vertex set V of the Cayley tree τ^k. In the sequel, we identify V with G_k.

The following assertion is quite obvious (see [2, 9]).

Theorem 2.1. *The group of left (right) shifts on the right (left) representation of the Cayley tree is the group of translations.*

By the group of translations we mean the automorphism group of the Cayley tree regarded as a graph. Recall (see, for example, [5]) that a mapping ψ on the vertex set of a graph G is called an automorphism of G if ψ preserves the adjacency relation, i.e., the images $\psi(u)$ and $\psi(v)$ of vertices u and v are adjacent if and only if u and v are adjacent.

For an arbitrary vertex $x_0 \in V$, we put

$$W_n = \left\{x \in V \mid d(x^0, x) = n\right\}, \quad V_n = \bigcup_{m=0}^{n} W_m, \quad L_n = \{l =< x, y >\in L \mid x, y \in V_n\}.$$

where $d(x, y)$ is the distance between x and y in the Cayley tree, i.e., the number of edges of the path between x and y.

For each $x \in G_k$, let $S(x)$ denote the set of immediate successor of x, i.e., if $x \in W_n$ then

$$S(x) = \{y \in W_{n+1} : d(x, y) = 1\}.$$

For each $x \in G_k$, let $S_1(x)$ denote the set of all neighbors of x, i.e., $S_1(x) = \{y \in G_k :< x, y >\in L\}$. The set $S_1(x) \setminus S(x)$ is a singleton. Let $x_\downarrow$ denote the (unique) element of this set.

Assume that spin takes its values in the set $\Phi = \{1, 2, \ldots, q\}$. By a configuration σ on V we mean a function taking $\sigma : x \in V \to \sigma(x) \in \Phi$. The set of all configurations coincides with the set $\Omega = \Phi^V$.

Consider the quotient group $G_k/G_k^* = \{H1, \ldots, H_r\}$, where G_k^* is a normal subgroup of index r with $r \geq 1$.

Definition 2.2. *A configuration $\sigma(x)$ is said to be G_k^*-periodic if $\sigma(x) = \sigma_i$ for all $x \in G_k$ with $x \in H_i$. A G_k-periodic configuration is said to be translation invariant.*

By *period* of a periodic configuration we mean the index of the corresponding normal subgroup.

Definition 2.3. *A configuration $\sigma(x)$ is said to be G_k^*-weakly periodic if $\sigma(x) = \sigma_{ij}$ for all $x \in G_k$ with $x_\downarrow \in H_i$ and $x \in H_j$.*

The Hamiltonian of the λ-model [7, 10] has a form,

$$H(\sigma) = \sum_{<x,y>\in L} \lambda(\sigma(x), \sigma(y)) \tag{1}$$

In what follows, we restrict ourself to the case $k = 2$ and $\Phi = \{1, 2, 3\}$, and for the sake of simplicity, we consider the following function:

$$\lambda(i,j) = \begin{cases} \overline{a} & if \quad |i-j| = 2, \\ \overline{b} & if \quad |i-j| = 1, \\ \overline{c} & if \quad i = j, \end{cases} \tag{2}$$

where $\overline{a}, \overline{b}, \overline{c} \in \mathbb{R}$ for some given numbers.

3. Ground States

In this section, we describe ground state of the λ-model on a Cayley tree. For a pair of configurations σ and φ coinciding almost everywhere, i.e., everywhere except finitely many points,we consider the relative Hamiltonian $H(\sigma, \varphi)$ determining the energy differences of the configurations σ and φ:

$$H(\sigma,\varphi) = \sum_{\substack{<x,y> \\ x,y\in V}} (\lambda(\sigma(x),\sigma(y)) - \lambda(\varphi(x),\varphi(y))) \tag{3}$$

Let M be the set of unit balls with vertices in V, i.e. $M = \{x, S_1(x), \forall x \in V\}$. We call the restriction of a configuration σ to the ball $b \in M$ a *bounded configuration* σ_b.

We define the energy of the configuration σ_b on b as follows

$$U(\sigma_b) = \frac{1}{2} \sum_{\substack{<x,y> \\ x,y\in V}} (\lambda(\sigma(x),\sigma(y)))$$

From (3), we got the following lemma.

We shall say that two bounded configurations σ_b and $\sigma'_{b'}$ belong to the same class if $U(\sigma_b) = U(\sigma'_{b'})$ and we write $\sigma'_{b'} \sim \sigma_b$.

Lemma 3.1. *Relative Hamiltonian* (3) *has the form*

$$H(\sigma,\varphi) = \sum_{b\in M} (U(\sigma_b) - U(\varphi_b)).$$

Lemma 3.2. *The inclusion*

$$U(\varphi_b) \in \left\{ \frac{\alpha+\beta+\gamma}{2}, \forall\ \alpha,\beta,\gamma \in \{\overline{a},\overline{b},\overline{c}\} \right\} \tag{4}$$

holds for every configuration φ_b on $b (b \in M)$.

Definition 3.3. *A configuration φ is called a ground state of the relative Hamiltonian H if*

$$U(\varphi_b) = \min \left\{ \frac{\alpha+\beta+\gamma}{2}, \forall\ \alpha,\beta.\gamma \in \{\overline{a},\overline{b},\overline{c}\} \right\} \tag{5}$$

for any $b \in M$

If a ground state is a periodic (resp. weakly periodic, translation invariant) configuration then we call it a periodic (resp. weakly periodic, translation invariant) ground state.

For any configuration σ_b, we have

$$U(\sigma_b) \in \{U_1, U_2, U_3, U_4, U_5, U_6, U_7, U_8, U_9, U_{10}\},$$

where

$$\begin{aligned} U_1 &= 3\overline{a}/2, & U_2 &= 3\overline{b}/2, \\ U_3 &= 3\overline{c}/2, & U_4 &= \overline{a}/2 + \overline{b}, \\ U_5 &= \overline{a}/2 + \overline{c}, & U_6 &= \overline{b}/2 + \overline{a}, \\ U_7 &= \overline{b}/2 + \overline{c}, & U_8 &= \overline{c}/2 + \overline{a}, \\ U_9 &= \overline{c}/2 + \overline{b}, & U_{10} &= (\overline{a} + \overline{b} + \overline{c})/2. \end{aligned} \tag{6}$$

We denote

$$A_m = \{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ U_m = \min_{1 \le k \le 10} \{U_k\}\} \tag{7}$$

Using (7), we obtain

$$\begin{aligned} A_1 &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{a} \le \overline{b} \le \overline{c}\right\} \cup \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{a} \le \overline{c} \le \overline{b}\right\}, \\ A_2 &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{b} \le \overline{c} \le \overline{a}\right\} \cup \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{b} \le \overline{a} \le \overline{c}\right\}, \\ A_3 &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{c} \le \overline{b} \le \overline{a}\right\} \cup \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{c} \le \overline{a} \le \overline{b}\right\}, \\ A_4 &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{a} = \overline{b} \le \overline{c}\right\}, \\ A_5 &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{a} = \overline{c} \le \overline{b}\right\}, \\ A_6 &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{a} = \overline{b} \le \overline{c}\right\}, \\ A_7 &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{b} = \overline{c} \le \overline{a}\right\}, \\ A_8 &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{a} = \overline{c} \le \overline{b}\right\}, \\ A_9 &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{b} = \overline{c} \le \overline{a}\right\}, \\ A_{10} &= \left\{(\overline{a}, \overline{b}, \overline{c}) \in \mathbb{R}^3 |\ \overline{a} = \overline{b} = \overline{c}\right\}, \mathbb{R}^3 = \cup_{i=1}^{10} A_i. \end{aligned}$$

Let us introduce some notations. We put

$$C_i = \{\sigma_b \in \Omega_b :\ U(\sigma_b) = u_i\}, \quad i = \overline{1, 10}$$

and $B^{(i)} = |\{x \in S_1(k) :\ \varphi_b(x) = i\}|$ for $i = \overline{1, 3}$.

Let $A \subset \{1, 2, ..., k+1\}$, $H_A = \{x \in G_k : \sum_{j \in A} w_j(x) - \text{even}\}$, where $w_j(x)$-is the number of letters a_j in the word x.

It is obvious, that H_A is a normal subgroup of index two [15]. Let $G_k/H_A = \{H_A, G_k \setminus H_A\}$ be the quotient group. We set $H_0 = H_A, H_1 = G_k \setminus H_A$.

4. Periodic Ground States

In this section, we describe H_0-periodic ground states. We note that each H_0-periodic configurations has the following form:

$$\sigma(x) = \begin{cases} \sigma_1, & \text{if} \quad x \in H_0, \\ \sigma_2, & \text{if} \quad x \in H_1 \end{cases} \tag{8}$$

where $\sigma_i \in \Phi = \{1, 2, 3\}$ $(i = 1, 2)$.

Theorem 4.1. *Let k=2 and $|A| = 1$. Then for the λ-model given by (2) the following statements hold:*

(i) *let $|\sigma_1 - \sigma_2| = 0$, then H_0-periodic ground state exists if and only if the parameters $\overline{a}, \overline{b}, \overline{c}$ belong to A_3;*

(ii) *If $|\sigma_1 - \sigma_2| = 1$, then H_0-periodic ground state exists if and only if the parameters $\overline{a}, \overline{b}, \overline{c}$ belong to A_7;*

(iii) *If $|\sigma_1 - \sigma_2| = 2$, then H_0-periodic ground state exists if and only if the parameters $\overline{a}, \overline{b}, \overline{c}$ belong to A_5.*

Proof. Let us consider the following configuration

$$\varphi(x) = \begin{cases} i, & \text{if} \quad x \in H_0 \\ i, & \text{if} \quad x \in H_1, \end{cases} \tag{9}$$

where $i = 1, 2, 3$.

By C_b we denote the center of $b \in M$. Let $C_b \in H_0$, then we have

$$\varphi_b(C_b) = i, \ B^{(i)} = 3;$$

Hence, $\varphi_b(x) \in C_3$, i.e. if $(\overline{a}, \overline{b}, \overline{c}) \in A_3$ then the corresponding configuration is a ground state.

Now consider the following configuration

$$\varphi(x) = \begin{cases} i, & \text{if} \quad x \in H_0, \\ j, & \text{if} \quad x \in H_1, \end{cases} \tag{10}$$

where $|i - j| = 1$

Assume that $C_b \in H_0$, the one finds

$$\varphi_b(C_b) = i, \ B^{(i)} = 2, \ B^{(j)} = 1;$$

Hence, $\varphi_b(x) \in C_7$.

Let $C_b \in H_1$, one has

$$\varphi_b(C_b) = j, \ B^{(i)} = 1, \ B^{(j)} = 2;$$

Hence, $\varphi_b(x) \in C_7$.

We conclude that, if $(\overline{a}, \overline{b}, \overline{c}) \in A_7$ then the corresponding periodic configuration $\varphi(x)$ is a H_0-periodic ground state.

Let us consider the following configuration

$$\varphi(x) = \begin{cases} i, & \text{if} \quad x \in H_0, \\ j, & \text{if} \quad x \in H_1, \end{cases} \tag{11}$$

where $|i - j| = 2$

Let $C_b \in H_0$, we have

$$\varphi_b(C_b) = i, \ B^{(i)} = 2, \ B^{(j)} = 1;$$

Hence, $\varphi_b(x) \in C_5$.

Let $C_b \in H_1$, one gets

$$\varphi_b(C_b) = j, \ B^{(i)} = 1, \ B^{(j)} = 2;$$

So, $\varphi_b(x) \in C_5$. Therefore, if $(\overline{a}, \overline{b}, \overline{c}) \in A_5$ then the corresponding periodic configuration $\varphi(x)$ is H_0-periodic ground state. This completes the proof. □

Remark 4.1. *Notice that periodic configurations (ground states) described in (i) Theorem 4.1 are translation-invariant. Full description of translation-invariant ground states has been carried out in* [10].

5. Weakly Periodic Ground States

In this section, we describe H_A-weakly periodic ground states. In what follows, H_A stands for the normal subgroup of index two. Due to the definition of weakly periodic configuration, we infer that each H_A-weakly periodic configuration has the following form:

$$\varphi(x)=\begin{cases}\sigma_1 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ \sigma_2 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ \sigma_3 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ \sigma_4 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases} \tag{12}$$

where $\sigma_i\in\Phi=\{1,2,3\}$, $(i=1,2,3,4)$.

Hence, all possible weakly periodic (non periodic) ground states have the following forms:

$$\varphi_1(x)=\begin{cases}1 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 1 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_2(x)=\begin{cases}1 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 1 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases}$$

$$\varphi_3(x)=\begin{cases}1 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 3 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_4(x)=\begin{cases}2 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 1 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 1 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 1 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases}$$

$$\varphi_5(x)=\begin{cases}2 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 1 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_6(x)=\begin{cases}2 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 1 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 3 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases}$$

$$\varphi_7(x)=\begin{cases}2 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 1 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_8(x)=\begin{cases}2 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 1 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases}$$

$$\varphi_9(x)=\begin{cases}2 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 3 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_{10}(x)=\begin{cases}2 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 3 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases}$$

$$\varphi_{11}(x)=\begin{cases}2 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 1 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 1 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_{12}(x)=\begin{cases}3 & if \quad x_\downarrow\in H_0, \quad x\in H_0,\\ 3 & if \quad x_\downarrow\in H_0, \quad x\in H_1,\\ 3 & if \quad x_\downarrow\in H_1, \quad x\in H_0,\\ 2 & if \quad x_\downarrow\in H_1, \quad x\in H_1,\end{cases}$$

$$\varphi_{13}(x)=\begin{cases}1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_{14}(x)=\begin{cases}1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1,\end{cases}$$

$$\varphi_{15}(x)=\begin{cases}1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_{16}(x)=\begin{cases}3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1,\end{cases}$$

$$\varphi_{17}(x)=\begin{cases}3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_{18}(x)=\begin{cases}3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1,\end{cases}$$

$$\varphi_{19}(x)=\begin{cases}3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1,\end{cases} \qquad \varphi_{20}(x)=\begin{cases}1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1,\end{cases}$$

Theorem 5.1. *Let $k=2$ and $|A|=1$. Then for the λ-model given by (2) the following statements hold:*

I. *There exist fourteen H_A-weakly periodic ground states if $(\overline{a},\overline{b},\overline{c})\in A_9$ that are not H_A-periodic, and they are described by φ_i, $i=1,\dots,14$. Moreover, they are not ground states on the set $\mathbb{R}^3\setminus A_9$.*

II. *There exist six H_A-weakly periodic ground states if $(\overline{a},\overline{b},\overline{c})\in A_5$ that are not H_A-periodic, and they are described by φ_i, $i=15,\dots,20$, and they are not ground states on the set $\mathbb{R}^3\setminus A_5$.*

III. *All H_A-weakly periodic (non periodic, in particular non translation-invariant) configurations (not mentioned in assertions I and II) are not weakly periodic ground states.*

Proof. 1) Consider

$$\varphi_1(x)=\begin{cases}1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, then we have the following possible cases

a) $C_{b_\downarrow}\in H_0$ and $\varphi_{1,b}(C_{b_\downarrow})=1$, then $\varphi_{1,b}(C_b)=1,\ B^{(1)}=3,\ \varphi_{1,b}\in C_3$,

b) $C_{b_\downarrow}\in H_0$ and $\varphi_{1,b}(C_{b_\downarrow})=2$, then $\varphi_{1,b}(C_b)=1,\ B^{(1)}=2,\ B^{(2)}=1,\ \varphi_{1,b}\in C_7$,

c) $C_{b_\downarrow}\in H_1$ and $\varphi_{1,b}(C_{b_\downarrow})=1$, this case is not exist, if $C_{b_\downarrow}\in H_1$, then $\varphi_{1,b}(C_{b_\downarrow})=2$,

d) $C_{b_\downarrow}\in H_1$ and $\varphi_{1,b}(C_{b_\downarrow})=2$, then $\varphi_{1,b}(C_b)=2,\ B^{(1)}=2,\ B^{(2)}=1,\ \varphi_{1,b}\in C_9$.

Let $C_b\in H_1$, then the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{1,b}(C_{b_\downarrow}) = 1$, then $\varphi_{1,b}(C_b) = 1,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{1,b} \in C_9$,
b) $C_{b_\downarrow} \in H_0$ and $\varphi_{1,b}(C_{b_\downarrow}) = 2$, this case is not exist, if $C_{b_\downarrow} \in H_0$, then $\varphi_{1,b}(C_{b_\downarrow}) = 1$,
c) $C_{b_\downarrow} \in H_1$ and $\varphi_{1,b}(C_{b_\downarrow}) = 1$, then $\varphi_{1,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{1,b} \in C_7$,
d) $C_{b_\downarrow} \in H_1$ and $\varphi_{1,b}(C_{b_\downarrow}) = 2$, then $\varphi_{1,b}(C_b) = 2,\ B^{(2)} = 3,\ \varphi_{1,b} \in C_3$.

Therefore, the configuration φ_1 is a weakly periodic ground state on the set $A_3 \cap A_7 \cap A_9 = A_7 = A_9$.

2) Consider

$$\varphi_2(x) = \begin{cases} 1 & if \quad x_\downarrow \in H_0, \quad x \in H_0, \\ 2 & if \quad x_\downarrow \in H_0, \quad x \in H_1, \\ 2 & if \quad x_\downarrow \in H_1, \quad x \in H_0, \\ 1 & if \quad x_\downarrow \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are
a) $C_{b_\downarrow} \in H_0$ and $\varphi_{2,b}(C_{b_\downarrow}) = 1$, then $\varphi_{2,b}(C_b) = 2,\ B^{(1)} = 2,\ B^{(2)} = 1,\ \varphi_{2,b} \in C_7$,
b) $C_{b_\downarrow} \in H_0$ and $\varphi_{2,b}(C_{b_\downarrow}) = 2$, then $\varphi_{2,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{2,b} \in C_9$,
c) $C_{b_\downarrow} \in H_1$ and $\varphi_{2,b}(C_{b_\downarrow}) = 1$, then $\varphi_{2,b}(C_b) = 2,\ B^{(1)} = 3,\ \varphi_{2,b} \in C_2$,
d) $C_{b_\downarrow} \in H_1$ and $\varphi_{2,b}(C_{b_\downarrow}) = 2$, this case is not exist, if $C_{b_\downarrow} \in H_1$, then $\varphi_{2,b}(C_{b_\downarrow}) = 1$.

Let $C_b \in H_1$, the possible cases are
a) $C_{b_\downarrow} \in H_0$ and $\varphi_{2,b}(C_{b_\downarrow}) = 1$, then $\varphi_{2,b}(C_b) = 2,\ B^{(1)} = 3,\ \varphi_{2,b} \in C_2$,
b) $C_{b_\downarrow} \in H_0$ and $\varphi_{2,b}(C_{b_\downarrow}) = 2$, this case is not exist, if $C_{b_\downarrow} \in H_0$, then $\varphi_{2,b}(C_{b_\downarrow}) = 1$,
c) $C_{b_\downarrow} \in H_1$ and $\varphi_{2,b}(C_{b_\downarrow}) = 1$, then $\varphi_{2,b}(C_b) = 1,\ B^{(1)} = 2,\ B^{(2)} = 1,\ \varphi_{2,b} \in C_7$,
d) $C_{b_\downarrow} \in H_1$ and $\varphi_{2,b}(C_{b_\downarrow}) = 2$, then $\varphi_{2,b}(C_b) = 1,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{2,b} \in C_9$.

Therefore, configuration φ_2 a weakly periodic ground state on the set $A_2 \cap A_7 \cap A_9 = A_7 = A_9$.

3) Consider

$$\varphi_3(x) = \begin{cases} 1 & if \quad x_\downarrow \in H_0, \quad x \in H_0, \\ 2 & if \quad x_\downarrow \in H_0, \quad x \in H_1, \\ 2 & if \quad x_\downarrow \in H_1, \quad x \in H_0, \\ 3 & if \quad x_\downarrow \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are
a) $C_{b_\downarrow} \in H_0$ and $\varphi_{3,b}(C_{b_\downarrow}) = 1$, then $\varphi_{3,b}(C_b) = 1,\ B^{(1)} = 2,\ B^{(2)} = 1,\ \varphi_{3,b} \in C_7$,
b) $C_{b_\downarrow} \in H_0$ and $\varphi_{3,b}(C_{b_\downarrow}) = 2$, then $\varphi_{3,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{3,b} \in C_9$,
c) $C_{b_\downarrow} \in H_1$ and $\varphi_{3,b}(C_{b_\downarrow}) = 2$, this case is not exist, if $C_{b_\downarrow} \in H_1$, then $\varphi_{3,b}(C_{b_\downarrow}) = 3$,

d) $C_{b_\downarrow} \in H_1$ and $\varphi_{3,b}(C_{b_\downarrow}) = 3$, then $\varphi_{3,b}(C_b) = 2,\ B^{(1)} = 2,\ B^{(3)} = 1,\ \varphi_{3,b} \in C_9$.

Let $C_b \in H_1$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{3,b}(C_{b_\downarrow}) = 1$, then $\varphi_{3,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(3)} = 2,\ \varphi_{3,b} \in C_2$,

b) $C_{b_\downarrow} \in H_0$ and $\varphi_{3,b}(C_{b_\downarrow}) = 2$, this case is not exist, if $C_{b_\downarrow} \in H_0$, then $\varphi_{3,b}(C_{b_\downarrow}) = 1$,

c) $C_{b_\downarrow} \in H_1$ and $\varphi_{3,b}(C_{b_\downarrow}) = 2$, then $\varphi_{3,b}(C_b) = 3,\ B^{(2)} = 2,\ B^{(3)} = 1,\ \varphi_{3,b} \in C_9$,

d) $C_{b_\downarrow} \in H_1$ and $\varphi_{3,b}(C_{b_\downarrow}) = 3$, then $\varphi_{3,b}(C_b) = 3,\ B^{(2)} = 1,\ B^{(3)} = 2,\ \varphi_{3,b} \in C_9$.

Therefore, configuration φ_3 a weakly periodic ground state on the set $A_2 \cap A_7 \cap A_9 = A_7 = A_9$.

4) consider

$$\varphi_4(x) = \begin{cases} 2 \quad if \quad x_\downarrow \in H_0, \quad x \in H_0, \\ 1 \quad if \quad x_\downarrow \in H_0, \quad x \in H_1, \\ 1 \quad if \quad x_\downarrow \in H_1, \quad x \in H_0, \\ 1 \quad if \quad x_\downarrow \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{4,b}(C_{b_\downarrow}) = 1$, then $\varphi_{4,b}(C_b) = 2,\ B^{(1)} = 2,\ B^{(2)} = 1,\ \varphi_{4,b} \in C_9$,

b) $C_{b_\downarrow} \in H_0$ and $\varphi_{4,b}(C_{b_\downarrow}) = 2$, then $\varphi_{4,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{4,b} \in C_7$,

c) $C_{b_\downarrow} \in H_1$ and $\varphi_{4,b}(C_{b_\downarrow}) = 1$, then $\varphi_{4,b}(C_b) = 1,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{4,b} \in C_9$.

Let $C_b \in H_1$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{4,b}(C_{b_\downarrow}) = 1$, then $\varphi_{4,b}(C_b) = 1,\ B^{(1)} = 3,\ \varphi_{4,b} \in C_3$,

b) $C_{b_\downarrow} \in H_0$ and $\varphi_{4,b}(C_{b_\downarrow}) = 2$, then $\varphi_{4,b}(C_b) = 1,\ B^{(1)} = 2,\ B^{(2)} = 1,\ \varphi_{4,b} \in C_7$,

c) $C_{b_\downarrow} \in H_1$ and $\varphi_{4,b}(C_{b_\downarrow}) = 1$, then $\varphi_{4,b}(C_b) = 1,\ B^{(1)} = 3,\ \varphi_{4,b} \in C_9$.

Therefore, configuration φ_4 a weakly periodic ground state on the set $A_3 \cap A_7 \cap A_9 = A_7 = A_9$.

5) Consider

$$\varphi_5(x) = \begin{cases} 2 \quad if \quad x_\downarrow \in H_0, \quad x \in H_0, \\ 1 \quad if \quad x_\downarrow \in H_0, \quad x \in H_1, \\ 2 \quad if \quad x_\downarrow \in H_1, \quad x \in H_0, \\ 2 \quad if \quad x_\downarrow \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{5,b}(C_{b_\downarrow}) = 2$, then $\varphi_{5,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{5,b} \in C_7$,

b) $C_{b_\downarrow} \in H_1$ and $\varphi_{5,b}(C_{b_\downarrow}) = 1$, this case is not exist, if $C_{b_\downarrow} \in H_1$, then $\varphi_{5,b}(C_{b_\downarrow}) = 2$,

c) $C_{b_\downarrow} \in H_1$ and $\varphi_{5,b}(C_{b_\downarrow}) = 2$, then $\varphi_{5,b}(C_b) = 2,\ B^{(2)} = 3,\ \varphi_{5,b} \in C_3$.

Let $C_b \in H_1$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{5,b}(C_{b_\downarrow}) = 2$, then $\varphi_{5,b}(C_b) = 1,\ B^{(2)} = 3,\ \varphi_{5,b} \in C_2$,

b) $C_{b_\downarrow} \in H_1$ and $\varphi_{5,b}(C_{b_\downarrow}) = 1$, then $\varphi_{5,b}(C_b) = 2,\ B^{(1)} = 2,\ B^{(2)} = 1,\ \varphi_{5,b} \in C_9$,

c) $C_{b_\downarrow} \in H_1$ and $\varphi_{5,b}(C_{b_\downarrow}) = 2$, then $\varphi_{5,b}(C_b) = 2,\ B^{(2)} = 3,\ \varphi_{5,b} \in C_3$.

Therefore, configuration φ_5 a weakly periodic ground state on the set $A_2 \cap A_3 \cap A_7 \cap A_9 = A_7 = A_9$.

6) Consider

$$\varphi_6(x) = \begin{cases} 2 & if \quad x_\downarrow \in H_0, \quad x \in H_0, \\ 1 & if \quad x_\downarrow \in H_0, \quad x \in H_1, \\ 3 & if \quad x_\downarrow \in H_1, \quad x \in H_0, \\ 2 & if \quad x_\downarrow \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{6,b}(C_{b_\downarrow}) = 2$, then $\varphi_{6,b}(C_b) = 2,\ B^{(2)} = 2,\ B^{(3)} = 3,\ \varphi_{6,b} \in C_7$,

b) $C_{b_\downarrow} \in H_0$ and $\varphi_{6,b}(C_{b_\downarrow}) = 3$, then $\varphi_{6,b}(C_b) = 2,\ B^{(1)} = B^{(2)} = B^{(3)} = 1,\ \varphi_{6,b} \in C_9$,

c) $C_{b_\downarrow} \in H_1$ and $\varphi_{6,b}(C_{b_\downarrow}) = 1$, this case is not exist, if $C_{b_\downarrow} \in H_1$, then $\varphi_{6,b}(C_{b_\downarrow}) = 2$,

d) $C_{b_\downarrow} \in H_1$ and $\varphi_{6,b}(C_{b_\downarrow}) = 2$, then $\varphi_{6,b}(C_b) = 3,\ B^{(2)} = 3,\ \varphi_{6,b} \in C_2$.

Let $C_b \in H_1$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{6,b}(C_{b_\downarrow}) = 2$, then $\varphi_{6,b}(C_b) = 1,\ B^{(2)} = 3,\ \varphi_{6,b} \in C_2$,

b) $C_{b_\downarrow} \in H_0$ and $\varphi_{6,b}(C_{b_\downarrow}) = 3$, this case is not exist, if $C_{b_\downarrow} \in H_0$, then $\varphi_{6,b}(C_{b_\downarrow}) = 2$,

c) $C_{b_\downarrow} \in H_1$ and $\varphi_{6,b}(C_{b_\downarrow}) = 1$, then $\varphi_{6,b}(C_b) = 2,\ B^{(1)} = B^{(2)} = B^{(3)} = 1,\ \varphi_{6,b} \in C_9$,

d) $C_{b_\downarrow} \in H_1$ and $\varphi_{6,b}(C_{b_\downarrow}) = 2$, then $\varphi_{6,b}(C_b) = 2,\ B^{(2)} = 2,\ B^{(3)} = 1,\ \varphi_{6,b} \in C_7$.

Therefore, configuration φ_6 a weakly periodic ground state on the set $A_2 \cap A_7 \cap A_9 = A_7 = A_9$.

7) Consider

$$\varphi_7(x) = \begin{cases} 2 & if \quad x_\downarrow \in H_0, \quad x \in H_0, \\ 2 & if \quad x_\downarrow \in H_0, \quad x \in H_1, \\ 1 & if \quad x_\downarrow \in H_1, \quad x \in H_0, \\ 2 & if \quad x_\downarrow \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{7,b}(C_{b_\downarrow}) = 1$, then $\varphi_{7,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{7,b} \in C_7$,

b) $C_{b_\downarrow} \in H_0$ and $\varphi_{7,b}(C_{b_\downarrow}) = 2$, then $\varphi_{7,b}(C_b) = 2,\ B^{(2)} = 3,\ \varphi_{7,b} \in C_3$,

c) $C_{b_\downarrow} \in H_1$ and $\varphi_{7,b}(C_{b_\downarrow}) = 2$, then $\varphi_{7,b}(C_b) = 1,\ B^{(2)} = 3,\ \varphi_{7,b} \in C_2$.

Let $C_b \in H_1$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{7,b}(C_{b_\downarrow}) = 1$, then $\varphi_{7,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{7,b} \in C_7$,

b) $C_{b_\downarrow} \in H_0$ and $\varphi_{7,b}(C_{b_\downarrow}) = 2$, then $\varphi_{7,b}(C_b) = 2,\ B^{(2)} = 3,\ \varphi_{7,b} \in C_3$,

c) $C_{b_\downarrow} \in H_1$ and $\varphi_{7,b}(C_{b_\downarrow}) = 2$, then $\varphi_{7,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{7,b} \in C_7$.

Therefore, configuration φ_7 a weakly periodic ground state on the set $A_2 \cap A_3 \cap A_7 = A_7$.

8) Consider

$$\varphi_8(x) = \begin{cases} 2 \quad if \quad x_\downarrow \in H_0, \quad x \in H_0, \\ 2 \quad if \quad x_\downarrow \in H_0, \quad x \in H_1, \\ 2 \quad if \quad x_\downarrow \in H_1, \quad x \in H_0, \\ 1 \quad if \quad x_\downarrow \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{8,b}(C_{b_\downarrow}) = 2$, then $\varphi_{8,b}(C_b) = 2,\ B^{(2)} = 3,\ \varphi_{8,b} \in C_3$,
b) $C_{b_\downarrow} \in H_1$ and $\varphi_{8,b}(C_{b_\downarrow}) = 1$, then $\varphi_{8,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{8,b} \in C_7$,
c) $C_{b_\downarrow} \in H_1$ and $\varphi_{8,b}(C_{b_\downarrow}) = 2$, this case is not exist, if $C_{b_\downarrow} \in H_1$, then $\varphi_{8,b}(C_{b_\downarrow}) = 1$.

Let $C_b \in H_1$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{8,b}(C_{b_\downarrow}) = 2$, then we can have two configuration:
   (i) $\varphi_{8,b}(C_b) = 2,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{8,b} \in C_7$,
   (ii) $\varphi_{8,b}(C_b) = 2,\ B^{(1)} = 3,\ \varphi_{8,b} \in C_1$,
b) $C_{b_\downarrow} \in H_1$ and $\varphi_{8,b}(C_{b_\downarrow}) = 1$, then $\varphi_{8,b}(C_b) = 1,\ B^{(1)} = 2,\ B^{(2)} = 1,\ \varphi_{8,b} \in C_7$,
c) $C_{b_\downarrow} \in H_1$ and $\varphi_{8,b}(C_{b_\downarrow}) = 2$, then $\varphi_{8,b}(C_b) = 1,\ B^{(1)} = 1,\ B^{(2)} = 2,\ \varphi_{8,b} \in C_9$.

Therefore, configuration φ_8 a weakly periodic ground state on the set $A_2 \cap A_3 \cap A_7 \cap A_9 = A_7 = A_9$.

9) Consider

$$\varphi_9(x) = \begin{cases} 2 \quad if \quad x_\downarrow \in H_0, \quad x \in H_0, \\ 2 \quad if \quad x_\downarrow \in H_0, \quad x \in H_1, \\ 2 \quad if \quad x_\downarrow \in H_1, \quad x \in H_0, \\ 3 \quad if \quad x_\downarrow \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{9,b}(C_{b_\downarrow}) = 2$, then $\varphi_{9,b}(C_b) = 2,\ B^{(2)} = 3,\ \varphi_{9,b} \in C_3$,
b) $C_{b_\downarrow} \in H_1$ and $\varphi_{9,b}(C_{b_\downarrow}) = 2$, this case is not exist, if $C_{b_\downarrow} \in H_1$, then $\varphi_{9,b}(C_{b_\downarrow}) = 3$,
c) $C_{b_\downarrow} \in H_1$ and $\varphi_{9,b}(C_{b_\downarrow}) = 3$, then $\varphi_{9,b}(C_b) = 2,\ B^{(2)} = 2,\ B^{(3)} = 1,\ \varphi_{9,b} \in C_7$.

Let $C_b \in H_1$, the possible cases are

a) $C_{b_\downarrow} \in H_0$ and $\varphi_{9,b}(C_{b_\downarrow}) = 2$, then $\varphi_{9,b}(C_b) = 2,\ B^{(2)} = 1,\ B^{(3)} = 2,\ \varphi_{9,b} \in C_9$,
b) $C_{b_\downarrow} \in H_1$ and $\varphi_{9,b}(C_{b_\downarrow}) = 2$, then $\varphi_{9,b}(C_b) = 3,\ B^{(2)} = 2,\ B^{(3)} = 1,\ \varphi_{9,b} \in C_9$,
c) $C_{b_\downarrow} \in H_1$ and $\varphi_{9,b}(C_{b_\downarrow}) = 3$, then $\varphi_{9,b}(C_b) = 3,\ B^{(2)} = 1,\ B^{(3)} = 2,\ \varphi_{9,b} \in C_7$.

Therefore, configuration φ_9 a weakly periodic ground state on the set $A_3 \cap A_7 \cap A_9 = A_7 = A_9$.

10) Consider

$$\varphi_{10}(x)=\begin{cases}2 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{10,b}(C_{b_\downarrow})=2$, then $\varphi_{10,b}(C_b)=2,\ B^{(2)}=3,\ \varphi_{10,b}\in C_3$,
b) $C_{b_\downarrow}\in H_0$ and $\varphi_{10,b}(C_{b_\downarrow})=3$, then $\varphi_{10,b}(C_b)=2,\ B^{(2)}=2,\ B^{(3)}=1,\ \varphi_{10,b}\in C_7$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{10,b}(C_{b_\downarrow})=2$, then $\varphi_{10,b}(C_b)=2,\ B^{(2)}=3,\ \varphi_{10,b}\in C_7$.

Let $C_b\in H_1$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{10,b}(C_{b_\downarrow})=2$, then $\varphi_{10,b}(C_b)=2,\ B^{(2)}=3,\ \varphi_{10,b}\in C_3$,
b) $C_{b_\downarrow}\in H_0$ and $\varphi_{10,b}(C_{b_\downarrow})=3$, this case is not exist, if $C_{b_\downarrow}\in H_0$, then $\varphi_{10,b}(C_{b_\downarrow})=2$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{10,b}(C_{b_\downarrow})=3$, then $\varphi_{10,b}(C_b)=2,\ B^{(2)}=2,\ B^{(3)}=1,\ \varphi_{10,b}\in C_7$.

Therefore, configuration φ_{10} a weakly periodic ground state on the set $A_2\cap A_3\cap A_7=A_7$.

11) Consider

$$\varphi_{11}(x)=\begin{cases}2 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{11,b}(C_{b_\downarrow})=1$, then $\varphi_{11,b}(C_b)=2,\ B^{(1)}=1,\ B^{(2)}=2,\ \varphi_{11,b}\in C_7$,
b) $C_{b_\downarrow}\in H_0$ and $\varphi_{11,b}(C_{b_\downarrow})=2$, then $\varphi_{11,b}(C_b)=2,\ B^{(2)}=3,\ \varphi_{11,b}\in C_3$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{11,b}(C_{b_\downarrow})=1$, then $\varphi_{11,b}(C_b)=1,\ B^{(1)}=1,\ B^{(2)}=2\ \varphi_{11,b}\in C_9$,
d) $C_{b_\downarrow}\in H_1$ and $\varphi_{11,b}(C_{b_\downarrow})=2$, then $\varphi_{11,b}(C_b)=1,\ B^{(1)}=2,\ B^{(2)}=1\ \varphi_{11,b}\in C_7$.

Let $C_b\in H_1$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{11,b}(C_{b_\downarrow})=1$, then $\varphi_{11,b}(C_b)=2,\ B^{(1)}=3,\ \varphi_{11,b}\in C_2$,
b) $C_{b_\downarrow}\in H_0$ and $\varphi_{11,b}(C_{b_\downarrow})=2$, then $\varphi_{11,b}(C_b)=2,\ B^{(1)}=2,\ B^{(2)}=1,\ \varphi_{11,b}\in C_9$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{11,b}(C_{b_\downarrow})=1$, then $\varphi_{11,b}(C_b)=1,\ B^{(1)}=3,\ \varphi_{11,b}\in C_3$,
d) $C_{b_\downarrow}\in H_1$ and $\varphi_{11,b}(C_{b_\downarrow})=2$, then $\varphi_{11,b}(C_b)=1,\ B^{(1)}=2,\ B^{(2)}=1,\ \varphi_{11,b}\in C_7$.

Therefore, configuration φ_{11} a weakly periodic ground state on the set $A_2\cap A_3\cap A_7\cap A_9=A_7=A_9$.

12) Consider

$$\varphi_{12}(x)=\begin{cases}3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{12,b}(C_{b_\downarrow})=3$, then $\varphi_{12,b}(C_b)=3,\ B^{(3)}=3,\ \varphi_{12,b}\in C_3$,
b) $C_{b_\downarrow}\in H_1$ and $\varphi_{12,b}(C_{b_\downarrow})=2$, then $\varphi_{12,b}(C_b)=3,\ B^{(2)}=1,\ B^{(3)}=2,\ \varphi_{12,b}\in C_7$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{12,b}(C_{b_\downarrow})=3$, this case is not exist, if $C_{b_\downarrow}\in H_1$, then $\varphi_{12,b}(C_{b_\downarrow})=2$.

Let $C_b\in H_1$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{12,b}(C_{b_\downarrow})=3$, then $\varphi_{12,b}(C_b)=3,\ B^{(2)}=2,\ B^{(3)}=1,\ \varphi_{12,b}\in C_9$,
b) $C_{b_\downarrow}\in H_1$ and $\varphi_{12,b}(C_{b_\downarrow})=2$, then $\varphi_{12,b}(C_b)=2,\ B^{(2)}=2,\ B^{(3)}=1,\ \varphi_{12,b}\in C_7$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{12,b}(C_{b_\downarrow})=3$, then $\varphi_{12,b}(C_b)=2,\ B^{(2)}=1,\ B^{(3)}=2,\ \varphi_{12,b}\in C_9$.

Therefore, configuration φ_{12} a weakly periodic ground state on the set $A_1\cap A_3\cap A_7\cap A_9=A_7=A_9$.

13) Consider

$$\varphi_{13}(x)=\begin{cases}1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{13,b}(C_{b_\downarrow})=1$, then $\varphi_{13,b}(C_b)=1,\ B^{(1)}=3,\ \varphi_{13,b}\in C_3$,
b) $C_{b_\downarrow}\in H_1$ and $\varphi_{13,b}(C_{b_\downarrow})=1$, this case is not exist, if $C_{b_\downarrow}\in H_1$, then $\varphi_{13,b}(C_{b_\downarrow})=2$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{13,b}(C_{b_\downarrow})=2$, then $\varphi_{3,b}(C_b)=2,\ B^{(1)}=2,\ B^{(2)}=1,\ \varphi_{13,b}\in C_7$.

Let $C_b\in H_1$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{13,b}(C_{b_\downarrow})=1$, then $\varphi_{13,b}(C_b)=1,\ B^{(1)}=1,\ B^{(2)}=2,\ \varphi_{13,b}\in C_9$,
b) $C_{b_\downarrow}\in H_1$ and $\varphi_{13,b}(C_{b_\downarrow})=1$, then $\varphi_{13,b}(C_b)=2,\ B^{(1)}=2,\ B^{(2)}=1,\ \varphi_{13,b}\in C_9$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{13,b}(C_{b_\downarrow})=2$, then $\varphi_{13,b}(C_b)=2,\ B^{(1)}=1,\ B^{(2)}=2,\ \varphi_{13,b}\in C_7$.

Therefore, configuration φ_{13} a weakly periodic ground state on the set $A_3\cap A_7\cap A_9=A_7=A_9$.

14) Consider

$$\varphi_{14}(x)=\begin{cases}1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 2 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{14,b}(C_{b_\downarrow})=1$, then $\varphi_{14,b}(C_b)=1,\ B^{(1)}=2,\ B^{(2)}=1,\ \varphi_{14,b}\in C_7$,
b) $C_{b_\downarrow}\in H_0$ and $\varphi_{14,b}(C_{b_\downarrow})=2$, then $\varphi_{14,b}(C_b)=1,\ B^{(1)}=1,\ B^{(2)}=2,\ \varphi_{14,b}\in C_9$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{14,b}(C_{b_\downarrow})=2$, then $\varphi_{14,b}(C_b)=1,\ B^{(1)}=1,\ B^{(2)}=2,\ \varphi_{14,b}\in C_9$.
Let $C_b\in H_1$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{14,b}(C_{b_\downarrow})=1$, then $\varphi_{14,b}(C_b)=2,\ B^{(1)}=1,\ B^{(2)}=2,\ \varphi_{14,b}\in C_7$,
b) $C_{b_\downarrow}\in H_1$ and $\varphi_{14,b}(C_{b_\downarrow})=2$, this case is not exist, if $C_{b_\downarrow}\in H_1$, then $\varphi_{14,b}(C_{b_\downarrow})=1$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{14,b}(C_{b_\downarrow})=2$, then $\varphi_{14,b}(C_b)=2,\ B^{(2)}=3,\ \varphi_{14,b}\in C_3$.
Therefore, configuration φ_{14} a weakly periodic ground state on the set $A_3\cap A_7\cap A_9=A_7=A_9$.

15) Consider

$$\varphi_{6}(x)=\begin{cases}1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{15,b}(C_{b_\downarrow})=1$, then $\varphi_{15,b}(C_b)=1,\ B^{(1)}=2,\ B^{(3)}=1,\ \varphi_{15,b}\in C_5$,
b) $C_{b_\downarrow}\in H_0$ and $\varphi_{15,b}(C_{b_\downarrow})=3$, then $\varphi_{15,b}(C_b)=1,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_{15,b}\in C_8$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{15,b}(C_{b_\downarrow})=3$, then $\varphi_{15,b}(C_b)=3,\ B^{(1)}=2,\ B^{(3)}=1,\ \varphi_{15,b}\in C_8$.
Let $C_b\in H_1$, the possible cases are
a) $C_{b_\downarrow}\in H_0$ and $\varphi_{15,b}(C_{b_\downarrow})=1$, then $\varphi_{15,b}(C_b)=3,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_{15,b}\in C_5$,
b) $C_{b_\downarrow}\in H_0$ and $\varphi_{15,b}(C_{b_\downarrow})=3$, this case is not exist, if $C_{b_\downarrow}\in H_0$, then $\varphi_{15,b}(C_{b_\downarrow})=1$,
c) $C_{b_\downarrow}\in H_1$ and $\varphi_{15,b}(C_{b_\downarrow})=3$, then $\varphi_{15,b}(C_b)=3,\ B^{(3)}=3,\ \varphi_{15,b}\in C_3$.
Therefore, configuration φ_{15} a weakly periodic ground state on the set $A_3\cap A_5\cap A_8=A_5=A_8$.

16) Consider

$$\varphi_{16}(x)=\begin{cases}3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, the possible cases are

a) $C_{b_\downarrow}\in H_0$ and $\varphi_{16,b}(C_{b_\downarrow})=1$, then $\varphi_{16,b}(C_b)=3,\ B^{(1)}=2,\ B^{(3)}=1,\ \varphi_{16,b}\in C_8$,

b) $C_{b_\downarrow}\in H_0$ and $\varphi_{16,b}(C_{b_\downarrow})=3$, then $\varphi_{16,b}(C_b)=3,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_{16,b}\in C_5$,

c) $C_{b_\downarrow}\in H_1$ and $\varphi_{16,b}(C_{b_\downarrow})=1$, this case is not exist, if $C_{b_\downarrow}\in H_1$, then $\varphi_{16,b}(C_{b_\downarrow})=3$,

d) $C_{b_\downarrow}\in H_1$ and $\varphi_{16,b}(C_{b_\downarrow})=3$, then $\varphi_{16,b}(C_b)=1,\ B^{(3)}=3,\ \varphi_{16,b}\in C_1$.

Let $C_b\in H_1$, the possible cases are

a) $C_{b_\downarrow}\in H_0$ and $\varphi_{16,b}(C_{b_\downarrow})=1$, then $\varphi_{16,b}(C_b)=1,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_{16,b}\in C_8$,

b) $C_{b_\downarrow}\in H_0$ and $\varphi_{16,b}(C_{b_\downarrow})=3$, then $\varphi_{16,b}(C_b)=1,\ B^{(3)}=3,\ \varphi_{16,b}\in C_2$,

c) $C_{b_\downarrow}\in H_1$ and $\varphi_{16,b}(C_{b_\downarrow})=1$, then $\varphi_{16,b}(C_b)=3,\ B^{(1)}=2,\ B^{(3)}=1,\ \varphi_{16,b}\in C_8$,

d) $C_{b_\downarrow}\in H_1$ and $\varphi_{16,b}(C_{b_\downarrow})=3$, then $\varphi_{16,b}(C_b)=3,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_{16,b}\in C_5$.

Therefore, configuration φ_{16} a weakly periodic ground state on the set $A_1\cap A_5\cap A_8=A_5=A_8$.

17) Consider

$$\varphi_{17}(x)=\begin{cases}3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, the possible cases are

a) $C_{b_\downarrow}\in H_0$ and $\varphi_{17,b}(C_{b_\downarrow})=1$, then $\varphi_{17,b}(C_b)=3,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_{17,b}\in C_5$,

b) $C_{b_\downarrow}\in H_0$ and $\varphi_{17,b}(C_{b_\downarrow})=3$, then $\varphi_{17,b}(C_b)=3,\ B^{(3)}=3,\ \varphi_{17,b}\in C_3$,

c) $C_{b_\downarrow}\in H_1$ and $\varphi_{17,b}(C_{b_\downarrow})=1$, then $\varphi_{17,b}(C_b)=1,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_{17,b}\in C_8$,

d) $C_{b_\downarrow}\in H_1$ and $\varphi_{17,b}(C_{b_\downarrow})=3$, this case is not exist, if $C_{b_\downarrow}\in H_1$, then $\varphi_{17,b}(C_{b_\downarrow})=1$.

Let $C_b\in H_1$, the possible cases are

a) $C_{b_\downarrow}\in H_0$ and $\varphi_{17,b}(C_{b_\downarrow})=1$, then $\varphi_{17,b}(C_b)=3,\ B^{(1)}=3,\ \varphi_{17,b}\in C_1$,

b) $C_{b_\downarrow}\in H_0$ and $\varphi_{17,b}(C_{b_\downarrow})=3$, then $\varphi_{17,b}(C_b)=3,\ B^{(1)}=2,\ B^{(3)}=1,\ \varphi_{17,b}\in C_8$,

c) $C_{b_\downarrow}\in H_1$ and $\varphi_{17,b}(C_{b_\downarrow})=1$, then $\varphi_{17,b}(C_b)=1,\ B^{(1)}=3,\ \varphi_{17,b}\in C_3$,

d) $C_{b_\downarrow}\in H_1$ and $\varphi_{17,b}(C_{b_\downarrow})=3$, then $\varphi_{17,b}(C_b)=1,\ B^{(1)}=2,\ B^{(3)}=1,\ \varphi_{17,b}\in C_5$.

Therefore, configuration φ_{17} a weakly periodic ground state on the set $A_1 \cap A_3 \cap A_5 \cap A_8 = A_5 = A_8$.

18) Consider

$$\varphi_{18}(x) = \begin{cases} 3 & if \quad x_{\downarrow} \in H_0, \quad x \in H_0, \\ 3 & if \quad x_{\downarrow} \in H_0, \quad x \in H_1, \\ 1 & if \quad x_{\downarrow} \in H_1, \quad x \in H_0, \\ 3 & if \quad x_{\downarrow} \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are

a) $C_{b_{\downarrow}} \in H_0$ and $\varphi_{18,b}(C_{b_{\downarrow}}) = 1$, then $\varphi_{18,b}(C_b) = 3$, $B^{(1)} = 1$, $B^{(3)} = 2$, $\varphi_{18,b} \in C_5$,

b) $C_{b_{\downarrow}} \in H_0$ and $\varphi_{18,b}(C_{b_{\downarrow}}) = 3$, then $\varphi_{18,b}(C_b) = 3$, $B^{(3)} = 3$, $\varphi_{18,b} \in C_3$,

c) $C_{b_{\downarrow}} \in H_1$ and $\varphi_{18,b}(C_{b_{\downarrow}}) = 3$, then $\varphi_{18,b}(C_b) = 1$, $B^{(3)} = 3$, $\varphi_{18,b} \in C_1$.

Let $C_b \in H_1$, the possible cases are

a) $C_{b_{\downarrow}} \in H_0$ and $\varphi_{18,b}(C_{b_{\downarrow}}) = 1$, then $\varphi_{18,b}(C_b) = 3$, $B^{(1)} = 1$, $B^{(3)} = 2$, $\varphi_{18,b} \in C_5$,

b) $C_{b_{\downarrow}} \in H_0$ and $\varphi_{18,b}(C_{b_{\downarrow}}) = 3$, then $\varphi_{18,b}(C_b) = 3$, $B^{(3)} = 3$, $\varphi_{18,b} \in C_3$,

c) $C_{b_{\downarrow}} \in H_1$ and $\varphi_{18,b}(C_{b_{\downarrow}}) = 3$, then $\varphi_{18,b}(C_b) = 3$, $B^{(1)} = 1$, $B^{(3)} = 2$, $\varphi_{18,b} \in C_5$.

Therefore, configuration φ_{18} a weakly periodic ground state on the set $A_1 \cap A_3 \cap A_5 = A_5$.

19) Consider

$$\varphi_{19}(x) = \begin{cases} 3 & if \quad x_{\downarrow} \in H_0, \quad x \in H_0, \\ 3 & if \quad x_{\downarrow} \in H_0, \quad x \in H_1, \\ 3 & if \quad x_{\downarrow} \in H_1, \quad x \in H_0, \\ 1 & if \quad x_{\downarrow} \in H_1, \quad x \in H_1. \end{cases}$$

Let $C_b \in H_0$, the possible cases are

a) $C_{b_{\downarrow}} \in H_0$ and $\varphi_{19,b}(C_{b_{\downarrow}}) = 3$, then $\varphi_{19,b}(C_b) = 3$, $B^{(3)} = 3$, $\varphi_{19,b} \in C_3$,

b) $C_{b_{\downarrow}} \in H_1$ and $\varphi_{19,b}(C_{b_{\downarrow}}) = 1$, then $\varphi_{19,b}(C_b) = 3$, $B^{(1)} = 1$, $B^{(3)} = 2$, $\varphi_{19,b} \in C_5$,

c) $C_{b_{\downarrow}} \in H_1$ and $\varphi_{19,b}(C_{b_{\downarrow}}) = 3$, then $\varphi_{19,b}(C_b) = 3$, $B^{(1)} = 1$, $B^{(3)} = 2$, $\varphi_{19,b} \in C_5$.

Let $C_b \in H_1$, the possible cases are

a) $C_{b_{\downarrow}} \in H_0$ and $\varphi_{19,b}(C_{b_{\downarrow}}) = 3$, then $\varphi_{19,b}(C_b) = 3$, $B^{(1)} = 2$, $B^{(3)} = 1$, $\varphi_{19,b} \in C_8$,

b) $C_{b_{\downarrow}} \in H_1$ and $\varphi_{19,b}(C_{b_{\downarrow}}) = 1$, then $\varphi_{19,b}(C_b) = 1$, $B^{(1)} = 2$, $B^{(3)} = 1$, $\varphi_{19,b} \in C_5$,

c) $C_{b_{\downarrow}} \in H_1$ and $\varphi_{19,b}(C_{b_{\downarrow}}) = 3$, then $\varphi_{19,b}(C_b) = 3$, $B^{(1)} = 1$, $B^{(3)} = 2$, $\varphi_{19,b} \in C_8$.

Therefore, configuration φ_{19} a weakly periodic ground state on the set $A_3 \cap A_5 \cap A_8 = A_5 = A_8$.

20) Consider

$$\varphi_{20}(x)=\begin{cases}1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

Let $C_b\in H_0$, the possible cases are

a) $C_{b_\downarrow}\in H_1$ and $\varphi_{20,b}(C_{b_\downarrow})=1$, then $\varphi_{20,b}(C_b)=3,\ B^{(1)}=3,\ \varphi_{20,b}\in C_3$,

b) $C_{b_\downarrow}\in H_0$ and $\varphi_{20,b}(C_{b_\downarrow})=1$, then $\varphi_{20,b}(C_b)=1,\ B^{(1)}=2,\ B^{(3)}=1,\ \varphi_{20,b}\in C_5$,

c) $C_{b_\downarrow}\in H_0$ and $\varphi_{20,b}(C_{b_\downarrow})=3$, then $\varphi_{20,b}(C_b)=1,\ B^{(1)}=2,\ B^{(3)}=1,\ \varphi_{20,b}\in C_5$.

Let $C_b\in H_1$, the possible cases are

a) $C_{b_\downarrow}\in H_0$ and $\varphi_{20,b}(C_{b_\downarrow})=1$, then $\varphi_{20,b}(C_b)=3,\ B^{(1)}=3,\ B^{(3)}=0,\ \varphi_{20,b}\in C_3$,

b) $C_{b_\downarrow}\in H_1$ and $\varphi_{20,b}(C_{b_\downarrow})=1$, then $\varphi_{20,b}(C_b)=3,\ B^{(1)}=2,\ B^{(3)}=1,\ \varphi_{20,b}\in C_8$,

c) $C_{b_\downarrow}\in H_1$ and $\varphi_{20,b}(C_{b_\downarrow})=3$, then $\varphi_{20,b}(C_b)=1,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_{20,b}\in C_5$.

Therefore, configuration φ_{20} a weakly periodic ground state on the set $A_3\cap A_5\cap A_8=A_5=A_8$.

Let we prove assertion III. We consider the remaining configurations. For example, let:

$$\varphi(x)=\begin{cases}3 & if \quad x_{\downarrow}\in H_0, \quad x\in H_0,\\ 1 & if \quad x_{\downarrow}\in H_0, \quad x\in H_1,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_0,\\ 3 & if \quad x_{\downarrow}\in H_1, \quad x\in H_1.\end{cases}$$

We show that $\varphi(x)$ is not a ground state.

Let $C_b\in H_0$, the possible cases are

a) $C_{b_\downarrow}\in H_0$ and $\varphi_b(C_{b_\downarrow})=3$, then $\varphi_b(C_b)=3,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_b\in C_8$,

b) $C_{b_\downarrow}\in H_1$ and $\varphi_b(C_{b_\downarrow})=3$, then $\varphi_b(C_b)=3,\ B^{(1)}=0,\ B^{(3)}=3,\ \varphi_b\in C_1$.

Let $C_b\in H_1$, the possible cases are

a) $C_{b_\downarrow}\in H_0$ and $\varphi_b(C_{b_\downarrow})=3$, then $\varphi_b(C_b)=1,\ B^{(1)}=0,\ B^{(3)}=3,\ \varphi_b\in C_3$,

b) $C_{b_\downarrow}\in H_1$ and $\varphi_b(C_{b_\downarrow})=1$, then $\varphi_b(C_b)=3,\ B^{(1)}=1,\ B^{(3)}=2,\ \varphi_b\in C_8$,

c) $C_{b_\downarrow}\in H_1$ and $\varphi_b(C_{b_\downarrow})=3$, then $\varphi_b(C_b)=3,\ B^{(1)}=0,\ B^{(3)}=3,\ \varphi_b\in C_1$.

We conclude that the configuration $\varphi(x)$ is a ground state on the set $\{(a,b,c)\in R^3: a=b=c\}$. Therefore, if $(a,b,c)\notin\{(a,b,c)\in R^3: a=b=c\}$, then the weakly periodic configuration $\varphi(x)$ is not a weakly periodic ground state.

As above, we analyze all possible cases, except for periodic (in particular translation-invariant) configurations and the configurations mentioned in assertions I and II we find that none of them is a ground state. This finishes the proof of Theorem 5.1. □

Remark 5.1. *Notice that the all configurations on the set* $\{(a,b,c)\in R^3: a=b=c\}$ *are ground state.*

Aknowledgments. The author RM thanks IIUM for providing financial support (grant FRGS 14-116-0357) and all facilities. The author also thanks the organizers of the 37-th International Conference on Quantum Probability and Related Topics (QP37) for travel support.

References

[1] G. I. Botirov, U. A. Rozikov, Potts model with competing interactions on the Cayley tree: The contour method, *Theoret. Math. Phys.* **153**(2007) 1423

[2] N. N. Ganikhodzhaev, Group representation and automorphisms of the Cayley tree, *Dokl. Akad. nauk Resp. Uzbekistan* **4** (1994) 3 [in Russian].

[3] N.N. Ganikhodjaev, F.M. Mukhamedov, J.F.F. Mendes, On the three state Potts model with competing interactions on the Bethe lattice, *Jour. Stat. Mech.* 2006, P08012

[4] M. I. Kargapolov, Yu. I. Merzlyakov, *Fundamentals of the Theory of Groups*, Springer-Verlag, Berlin, 1979.

[5] O. Melnikov, R. I. Tyshkevich, V. A. Yemelichev, V. I. Sarvanov, *Lectures on Graph Theory* B. I. Wissenschaftsverlag, Mannheim ,1994

[6] R. A. Minlos, *Introduction to Mathematical Statistical Physics*, Amer. Math. Soc., Providence, RI, 2000.

[7] F.M. Mukhamedov, On a factor associated with the unordered phase of λ-model on a Cayley tree. *Rep. Math. Phys.* **53** (2004), 1–18.

[8] F.M.Mukhamedov, U.A.Rozikov, Extremality of the disordered phase of the nonhomogeneous Potts model on the Cayley tree. *Theor. Math. Phys.* **124** (2000), 1202–1210

[9] F.M. Mukhamedov, U.A. Rozikov U.A., J.F.F. Mendes, On contour arguments for the three state Potts model with competing interactions on a semi-infinite Cayley tree, *Jour. Math. Phys.* **48** (2007) 013301,

[10] F. Mukhamedov, C. H. Pah, H. Jamil, On Ground States and Phase Transitions of λ-Model on the Cayley Tree, `arXiv:1610.07019`

[11] M. M. Rakhmatullaev, Description of weak periodic ground states of Ising model with competing interactions on Cayley tree, *Appl. Math. Inf. Sci.* **4**(2010) 237.

[12] M. M. Rakhmatullaev, Weakly periodic Gibbs measures and ground states for the Potts model with competing interactions on the Cayley tree, *Theoret. Math. Phys.* **176**(2013) 1236

[13] M.M. Rahmatullaev, M.A. Rasulova, Periodic and Weakly Periodic Ground States for the Potts Model with Competing Interactions on the Cayley Tree, *Siberian Adv. Math.*, **26**(2016) 215229

[14] U.A. Rozikov, Description of limit Gibbs measures for λ-models on the Bethe lattice, *Siberan Math. Jour.* **39**(1998), 373380

[15] U. A. Rozikov, *Gibbs Measures on Cayley Trees,* World Scientific, Hackensack, 2013

[16] U.A. Rozikov, R.M. Khakimov, Periodic Gibbs measures for the Potts model on the Cayley tree, *Theor. Math. Phys.* **175**(2013), 699709.

[17] U. A. Rozikov and M. M. Rakhmatullaev, Weakly periodic ground states and Gibbs measures for the Ising model with competing interactions on the Cayley tree, *Theoret. Math. Phys.* **160** (2009) 1292

[18] Ya.G.Sinai, *Theory of Phase Transitions:Rigorous Results*, Pergamon Press, Oxford, 1982

Farrukh Mukhamedov, Department of Mathematical Sciences, College of Science, The United Arab Emirates University, P.O. Box, 15551, Al Ain, Abu Dhabi, UAE
E-mail address: far75m@gmail.com farrukh.m@uaeu.ac.ae

Chin Hee Pah, Department of Computational & Theoretical Sciences,, Faculty of Science, International Islamic University Malaysia,, P.O. Box, 141, 25710, Kuantan, Pahang, Malaysia
E-mail address: pahchinhee@iium.edu.my

Muzaffar Rahmatullaev, Institute of Mathematics, National University of Uzbekistan, 29, Do'rmon Yo'li str., 100125, Tashkent, Uzbekistan.
E-mail address: mrahmatullaev@rambler.ru

Hakim Jamil, Department of Computational & Theoretical Sciences,, Faculty of Science, International Islamic University Malaysia,, P.O. Box, 141, 25710, Kuantan, Pahang, Malaysia
E-mail address: m.hakimjamil@yahoo.com.my